

RIESZ BASES OF REPRODUCING KERNELS IN FOCK TYPE SPACES AND DE BRANGES SPACES

ANTON BARANOV, YURII BELOV, ALEXANDER BORICHEV

ABSTRACT. We describe the radial Fock type spaces which possess Riesz bases of normalized reproducing kernels and which are (are not) isomorphic to de Branges spaces in terms of the weight functions.

1. INTRODUCTION AND MAIN RESULTS

Let \mathcal{F} be a Hilbert space of entire functions. Assume that (i) \mathcal{F} possesses the division property, that is, if $f \in \mathcal{F}$, $f(\lambda) = 0$, then $f/(\cdot - \lambda) \in \mathcal{F}$, and (ii) \mathcal{F} possesses the bounded point evaluation property, that is, for each $\lambda \in \mathbb{C}$, the mapping $L_\lambda : f \mapsto f(\lambda)$ is a bounded linear functional on \mathcal{F} . For every $\lambda \in \mathbb{C}$ there exists $\mathbf{k}_\lambda \in \mathcal{F}$, the reproducing kernel at λ in \mathcal{F} :

$$f(\lambda) = \langle f, \mathbf{k}_\lambda \rangle_{\mathcal{F}}, \quad f \in \mathcal{F}.$$

Let $\mathbb{k}_\lambda = \mathbf{k}_\lambda / \|\mathbf{k}_\lambda\|$ be the normalized reproducing kernel at λ . Given a sequence $\Lambda \subset \mathbb{C}$, we say that $\{\mathbb{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis (of normalized reproducing kernels) in \mathcal{F} if it is complete and for some $c, C > 0$ we have

$$c \sum_{\lambda \in \Lambda} |a_\lambda|^2 \leq \left\| \sum_{\lambda \in \Lambda} a_\lambda \mathbb{k}_\lambda \right\|^2 \leq C \sum_{\lambda \in \Lambda} |a_\lambda|^2,$$

for finite sequences $\{a_\lambda\}$ of complex numbers. Equivalently, $\{\mathbb{k}_\lambda\}_{\lambda \in \Lambda}$ is a linear isomorphic image of an orthonormal basis in a separable Hilbert space.

The de Branges spaces $\mathcal{H}(\mathcal{E})$ are Hilbert spaces of entire functions determined by Hermite–Biehler class entire functions \mathcal{E} , see [5]. The norm in these spaces is given by $\|F\|_{\mathcal{H}(\mathcal{E})}^2 = \int_{\mathbb{R}} |F(x)|^2 / |\mathcal{E}(x)|^2 dx$. They possess Riesz bases of normalized reproducing kernels at real points $\{t_n\}_{n \in \mathbb{N}}$. Correspondingly, every space $\mathcal{H}(\mathcal{E})$ can be identified with the space \mathcal{H} of all entire functions of the form

$$F(z) = A(z) \sum_{n \in \mathbb{N}} \frac{a_n \mu_n^{1/2}}{z - t_n}, \quad \{a_n\} \in \ell^2,$$

where $\{t_n\}_{n \in \mathbb{N}}$ is an increasing sequence such that $|t_n| \rightarrow \infty$, $|n| \rightarrow \infty$, $N = \mathbb{Z}$, \mathbb{Z}_+ or \mathbb{Z}_- , $\sum_{n \in \mathbb{N}} \mu_n \delta_{t_n}$ is a positive measure on \mathbb{R} satisfying $\sum_{n \in \mathbb{N}} \frac{\mu_n}{t_n^2 + 1} < \infty$, A is an entire function with zero set $\{t_n\}_{n \in \mathbb{N}}$ which is real on the real line, and the norm of F is defined as $\|F\|_{\mathcal{H}} = \|\{a_n\}\|_{\ell^2}$. On the other hand, every \mathcal{F} as above possessing a Riesz basis of normalized reproducing kernels at real points is (up to the norm equivalence) a de Branges space; see, for instance, [2].

The work was supported by Russian Science Foundation grant 14-41-00010.

Given a continuous function (a weight) h defined on $[0, +\infty)$, we extend it to the whole complex plane \mathbb{C} by $h(z) = h(|z|)$, and consider the Fock type space

$$\mathcal{F}_h = \{f \in \text{Hol}(\mathbb{C}) : \|f\|^2 = \|f\|_h^2 = \int_{\mathbb{C}} |f(z)|^2 e^{-h(z)} dm(z) < \infty\},$$

where dm is area Lebesgue measure. This is a Hilbert space of entire functions satisfying properties (i) and (ii).

Seip proved in [10] that the standard Fock space \mathcal{F}_h , $h(z) = |z|^2$, has no Riesz basis of normalized reproducing kernels. Later on, it was proved that \mathcal{F}_h has no Riesz basis of normalized reproducing kernels for sufficiently regular h such that $h(t) \gg (\log t)^2$ [3, 4]. On the other hand, the spaces \mathcal{F}_h with $h(t) = (\log t)^\alpha$, $1 < \alpha \leq 2$, possess Riesz bases of normalized reproducing kernels at real points [4], and hence, are de Branges spaces.

In the opposite direction, Theorem 1.2 in [1] states that the de Branges space \mathcal{H} determined by the spectral data $(\{t_n\}_{n \in \mathbb{N}}, \{\mu_n\}_{n \in \mathbb{N}})$ coincides (up to the norm equivalence) with a Fock type space \mathcal{F}_h if and only if (i) the sequence $\{t_n\}_{n \in \mathbb{N}}$ is lacunary: $\liminf_{t_n \rightarrow \infty} \frac{t_{n+1}}{t_n} > 1$, $\liminf_{t_n \rightarrow -\infty} \frac{|t_n|}{|t_{n+1}|} > 1$, and (ii) for some $C > 0$ and any n ,

$$\sum_{|t_k| \leq |t_n|} \mu_k + t_n^2 \sum_{|t_k| > |t_n|} \frac{\mu_k}{t_k^2} \leq C \mu_n.$$

(These de Branges spaces \mathcal{H} are characterized by the property that every complete and minimal system of reproducing kernels is a strong Markushevich basis (strong M -basis), see [1].)

The main question we deal with in this paper is to find conditions on h determining whether the Fock type space \mathcal{F}_h has a Riesz basis of (normalized) reproducing kernels and whether \mathcal{F}_h is a de Branges space, or, correspondingly, whether the norm in \mathcal{F}_h is equivalent to a weighted L^2 norm along the real line. Our work can be considered as completing a series of works [4], [2], and [1].

From now on let us fix $\psi(t) = h(e^t)$.

Theorem 1.1. *If ψ'' increases to $+\infty$, then \mathcal{F}_h has no Riesz bases of (normalized) reproducing kernels.*

In this case, $(\log t)^2 = o(h(t))$, $t \rightarrow \infty$. Under different (in some cases more restrictive) regularity conditions, this result follows from [10], [3], and [4].

Theorem 1.2. *If $\lim_{t \rightarrow \infty} \psi'(t) = \infty$, ψ'' is a non-increasing positive function, and $|\psi'''(t)| = O(\psi''(t)^{5/3})$, $t \rightarrow \infty$, then \mathcal{F}_h has a Riesz basis of (normalized) reproducing kernels at real points, and hence, is (up to the norm equivalence) a de Branges space.*

Theorem 1.3. *Given any convex positive function φ with $x = o(\varphi(x))$, $x \rightarrow \infty$, there exists a function ψ increasing to $+\infty$ and satisfying the property $\psi(t) = o(\varphi(t))$, $t \rightarrow \infty$, such that \mathcal{F}_h has a Riesz basis of (normalized) reproducing kernels and is not (up to the norm equivalence) a de Branges space.*

1.1. Let $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ be a complete minimal family of the reproducing kernels in \mathcal{F}_h . Then there exists a so called generating function E with simple zeros at the points of Λ such that $E \notin \mathcal{F}_h$ and $E_\lambda = E/(\cdot - \lambda) \in \mathcal{F}_h$ for every $\lambda \in \Lambda$. The family $\{\|\mathbf{k}_\lambda\|E_\lambda/E'(\lambda)\}_{\lambda \in \Lambda}$ is biorthogonal to $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$. Consider the mappings $T_\Lambda : f \mapsto f|_\Lambda$ and

$$R_\Lambda : \{a_\lambda\}_{\lambda \in \Lambda} \mapsto E(z) \sum_{\lambda \in \Lambda} \frac{a_\lambda}{E'(\lambda)} \cdot \frac{1}{z - \lambda}.$$

for finite sequences $\{a_\lambda\}_{\lambda \in \Lambda}$. Then $T_\Lambda R_\Lambda = Id$. If T_Λ is bounded from \mathcal{F}_h to $\ell^2(1/\|\mathbf{k}_\lambda\|^2)$ and R_Λ is bounded from $\ell^2(1/\|\mathbf{k}_\lambda\|^2)$ to \mathcal{F}_h , then the family $\{\|\mathbf{k}_\lambda\|E_\lambda/E'(\lambda)\}_{\lambda \in \Lambda}$ is a Riesz basis, and hence $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis of normalized reproducing kernels in \mathcal{F}_h (see, for instance, [9, Section C.3.1]).

1.2. In this paper, we deal mainly with non-decreasing weights h . This does restrict the class of Fock type spaces we consider.

Proposition 1.4. *There exist continuous h_0 such that \mathcal{F}_{h_0} does not coincide (up to the norm equivalence) with \mathcal{F}_h for any non-decreasing weight h .*

Let us introduce some notation. If ψ is convex, then h is subharmonic and we set $\rho(z) = (\Delta h(z))^{-1/2}$, $\tau(t) = (\psi''(t))^{-1/2}(t) = \rho(e^t)e^{-t}$. Later on, we'll see that ρ will be a local metric coefficient and τ will be a local metric coefficient in the logarithmic scale. Given an analytic function f , we denote by $Z(f)$ its zero set.

The proofs of Theorems 1.1–1.3 are contained in Sections 2–4, correspondingly. Proposition 1.4 is proved in Section 5.

2. PROOF OF THEOREM 1.1

We follow the method proposed in [10] and later used in [4]. It turns out that in some large annuli we can obtain precise estimates on the norm of the reproducing kernel. If $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis, then Λ is ρ -separated and ρ -dense on these annuli (see Lemma 2.6 below) and this implies that the corresponding Hilbert transform is unbounded which leads to a contradiction.

We suppose in this section that ψ'' increases to $+\infty$, τ decreases to 0.

Lemma 2.1. *Given $A < \infty$, $y_0 > 0$, there exists $y \geq y_0$ such that*

$$(2.1) \quad \left| \frac{\psi''(x)}{\psi''(y)} - 1 \right| \leq \frac{1}{A},$$

if $|x - y| \leq A\tau(y)$.

For similar statements see, for instance, [7].

Proof. Without loss of generality, A is sufficiently large. Set

$$y_n = y_{n-1} + 4A\tau(y_{n-1}), \quad n \geq 1,$$

and suppose that

$$\psi''(y_n) \geq \frac{A+1}{A} \psi''(y_{n-1}), \quad n \geq 1.$$

Then

$$\psi''(y_n) \geq \left(\frac{A+1}{A}\right)^n \psi''(y_0), \quad n \geq 1,$$

and, hence,

$$\sum_{n \geq 1} (y_n - y_{n-1}) < \infty, \\ \lim_{n \rightarrow \infty} y_n = y < \infty,$$

which is absurd.

Hence, there exists $z \geq y_0$ such that

$$\psi''(z) \leq \psi''(x) < \frac{A+1}{A} \psi''(z), \quad 0 \leq x - z \leq 4A\tau(z).$$

It remains to set $y = z + 2A\tau(z)$. For large A we obtain (2.1). \square

Lemma 2.2. *Let $B > 0$, $N < \infty$. Given $e^y = R_1 < R_2 < \dots < R_N$, $R_{N+1} = +\infty$, and integers k_1, \dots, k_N such that*

$$\frac{1}{B} \leq \frac{R_{j+1} - R_j}{e^{y\tau(y)}} \leq B, \quad 1 \leq j < N, \\ \frac{1}{B} \leq k_j \tau(y) \leq B, \quad 1 \leq j \leq N,$$

we set

$$P(z) = \prod_{1 \leq j \leq N} \left(1 - \left(\frac{z}{R_j}\right)^{k_j}\right).$$

Next, we set

$$v(t) = \begin{cases} 0, & t < R_1, \\ \sum_{1 \leq s \leq j} k_s (\log t - \log R_s), & R_j \leq t < R_{j+1}, 1 \leq j \leq N. \end{cases}$$

Then for every $\delta > 0$ for some $D = D(\delta, B)$ independent of N and for all $y > y(\delta, \psi, B)$ we have

$$|\log |P(z)| - v(|z|)| \leq D(\delta, B)$$

if $\text{dist}(z, Z(P)) > \delta e^y \tau(y)$.

Proof. It suffices to use that

$$|\log |1 - z^k|| \leq C e^{-t}, \quad |z| = 1 - \frac{t}{k}, \quad t \geq 1, \\ |\log |1 - z^k|| \leq D_0(\delta), \quad 1 - \frac{1}{k} < |z| < 1 + \frac{1}{k}, \quad \text{dist}(z, \Lambda_k) > \frac{\delta}{k}, \\ |\log |1 - z^k| - k \log |z|| \leq C e^{-c \min(t, k)}, \quad |z| = 1 + \frac{t}{k}, \quad t \geq 1,$$

where $\Lambda_k = \{e^{2\pi i j/k}\}_{0 \leq j < k}$. \square

Given $A, R > 0$, set

$$\Omega_{R,A} = \{z \in \mathbb{C} : ||z| - R| \leq A\rho(R)\}.$$

Lemma 2.3. *Given $A < \infty$, there exist $R > 0$ and a polynomial Q such that $Z(Q) \setminus \{0\} \subset \Omega_{R,2A}$ and*

$$\begin{aligned} \frac{1}{M} &\leq \frac{|Q(z)|\rho(R)e^{-h(z)}}{\text{dist}(z, Z(Q))} \leq M, & z \in \Omega_{R,A}, \\ |Q(z)| &\leq Me^{h(z)}, & z \in \mathbb{C}, \\ \text{dist}(w, Z(Q) \setminus \{w\}) &\geq \frac{\rho(R)}{M}, & w \in Z(Q) \setminus \{0\}, \\ \text{dist}(z, Z(Q) \setminus \{w\}) &\leq M\rho(R), & z \in \Omega_{R,A}, \end{aligned}$$

for some absolute constant M .

Proof. The lemma follows immediately from Lemmas 2.1 and 2.2. Indeed, by Lemma 2.1 we can find R such that ψ'' is almost constant on the corresponding interval around $\log R$. Then we find k_j, R_j in Lemma 2.2 and $a \in \mathbb{R}, k \in \mathbb{N}$ such that $a + kt + v(\exp t)$ is close to $\psi(t)$ for $\exp t \in \Omega_{R,2A}$. Finally, we set $Q(z) = e^a z^k P(z)$, where P is given by Lemma 2.2. \square

For a similar construction see [3]; for an alternative way to produce analytic functions Q satisfying such conditions see the atomization procedure in [8].

Now we can proceed like in [4].

Lemma 2.4. *In the conditions of Lemma 2.3, given $w \in \Omega_{R,A/2}$, there exists a function Φ_w analytic in the disc $D_w = \{z \in \mathbb{C} : |z - w| < \rho(R)\}$ and such that*

$$\frac{1}{M} \leq |\Phi_w(z)|e^{-h(z)} \leq M, \quad z \in D_w,$$

for some absolute constant M .

Proof. The proof of this folklore result is identical to that of Lemma 2.1 in [4]. \square

Lemma 2.5. *In the conditions of Lemma 2.3,*

$$\frac{1}{M} \leq \|\mathbf{k}_z\|e^{-h(z)/2}\rho(R) \leq M, \quad z \in \Omega_{R,A/3},$$

for some absolute constant M .

Proof. Analogous to that of Lemma 2.3 in [4] with F there replaced by our polynomial Q . \square

Lemma 2.6. *In the conditions of Lemma 2.3, let a sequence $\Lambda \subset \mathbb{C}$ be such that $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis in \mathcal{F}_h . Then*

- (a) $\text{dist}(\lambda, \Lambda \setminus \{\lambda\}) \geq \beta\rho(R), \quad \lambda \in \Lambda \cap \Omega_{R,A/4},$
- (b) $\text{dist}(z, \Lambda) \leq \rho(R)/\beta, \quad z \in \Omega_{R,A/4},$

for some $\beta = \beta(\Lambda)$ and for $A > A(\Lambda)$.

Proof. Analogous to that of Lemma 2.4 in [4]. \square

Proof of Theorem 1.1. We follow the proof of Theorem 2.5 in [4]. Suppose that the system $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis in \mathcal{F}_h . Fix large A , choose R and

Q as in Lemma 2.3, and define E as in Subsection 1.1. Lemma 2.5 implies that

$$\left\| \frac{E}{\cdot - \lambda} \right\|^2 \asymp \frac{|E'(\lambda)|^2}{\|\mathbf{k}_z\|^2} \asymp |E'(\lambda)|^2 \rho^2(\lambda) e^{-h(\lambda)}, \quad \lambda \in \Lambda \cap \Omega_{R, A/4}.$$

Consider the function $E/[(\cdot - \lambda)\Phi_\lambda]$. Applying the mean value property we obtain that

$$\int_{\mathbb{C}} \frac{|E(z)|^2}{|z - \lambda|^2} e^{-h(z)} dm(z) \leq \frac{C}{\rho^2(R)} \int_{|z - \lambda| < \rho(R)} |E(z)|^2 e^{-h(z)} dm(z).$$

By Lemma 2.6 (a), we have

$$\begin{aligned} & \int_{\mathbb{C}} \left(\sum_{|\lambda - R| < N\rho(R), \lambda \in \Lambda} \frac{1}{|z - \lambda|^2} \right) |E(z)|^2 e^{-h(z)} dm(z) \\ & \leq \frac{C}{\rho^2(R)} \int_{|R - z| < (N+1)\rho(R)} |E(z)|^2 e^{-h(z)} dm(z), \end{aligned}$$

with $A/4 - 1 < N \leq A/4$, and, as a result,

$$(2.2) \quad \inf_{z: |z - R| < (N+1)\rho(R)} \left[\rho^2(R) \left(\sum_{|\lambda - R| < N\rho(R), \lambda \in \Lambda} \frac{1}{|z - \lambda|^2} \right) \right] \leq C.$$

Finally, by Lemma 2.6 (b),

$$\rho^2(R) \left(\sum_{|\lambda - R| < N\rho(R), \lambda \in \Lambda} \frac{1}{|z - \lambda|^2} \right) \geq C \int_{\rho(R) < |\zeta| < N\rho(R)} \frac{dm(\zeta)}{|z - \zeta|^2}.$$

For large A (and hence, large N) we get a contradiction to (2.2). \square

3. PROOF OF THEOREM 1.2

The situation for h of slow growth is quite different from that considered in Section 2. In particular, along some special (lacunary) sequence of points $\{\lambda_n\}_{n \geq 0}$ we obtain here that $\|\mathbf{k}_{\lambda_n}\|^2 \asymp e^{h(\lambda_n)}/(\lambda_n \rho(\lambda_n)) \gtrsim e^{h(\lambda_n)}/(\rho(\lambda_n))^2$ instead of $\|\mathbf{k}_{\lambda_n}\|^2 \asymp e^{h(\lambda_n)}/(\rho(\lambda_n))^2$ in Lemma 2.5. To prove the boundedness of the operator R_Λ introduced in Subsection 1.1 we use the results of [2].

For $n \geq \psi'(0)/2 - 1$ we choose y_n such that $\psi'(y_n) = 2n + 2$ and set $\lambda_n = \exp y_n$. Define

$$g(t) = \psi(y_n + t) - (y_n + t)\psi'(y_n), \quad -y_n \leq t < \infty.$$

Set $\alpha_n = \psi''(y_n) = g''(0)$. Then $g'(0) = 0$, $0 < g''(t) \leq \psi''(0)$, $|g'''(t)| = O(g''(t)^{5/3})$, and hence,

$$(3.1) \quad g(t) - g(0) \asymp \alpha_n t^2, \quad |t| \leq C\alpha_n^{-2/3},$$

$$(3.2) \quad g'(t) \asymp \alpha_n t, \quad |t| \leq C\alpha_n^{-2/3},$$

$$(3.3) \quad g(t) - g(0) \geq C\alpha_n^{-1/3} + C\alpha_n^{1/3}|t|, \quad |t| \geq C\alpha_n^{-2/3}.$$

We assign to the finite number of y_n not yet defined, arbitrary values in such a way that the sequence $\{y_n\}_{n \geq 0}$ of positive numbers is strictly increasing.

Lemma 3.1.

$$\|z^n\|^2 \asymp e^{(2n+2)y_n - \psi(y_n)} (\psi''(y_n))^{-1/2}, \quad n \geq 0.$$

Proof. We have

$$\|z^n\|^2 = O(1) + 2\pi \int_1^\infty t^{2n+1} e^{-\psi(\log t)} dt = O(1) + 2\pi \int_0^\infty e^{(2n+2)s - \psi(s)} ds.$$

By (3.1) and (3.3) we have

$$\frac{C_1}{\sqrt{\alpha_n}} \leq \int_{-y_n}^\infty e^{g(0) - g(t)} dt \leq \frac{C_2}{\sqrt{\alpha_n}}$$

and hence,

$$\frac{C_1}{\sqrt{\alpha_n}} \leq \|z^n\|^2 e^{\psi(y_n) - (2n+2)y_n} \leq \frac{C_2}{\sqrt{\alpha_n}}.$$

□

Next, we estimate the norm of the reproducing kernel at the points λ_n ; compare also to Lemma 2.5.

Lemma 3.2.

$$\|\mathbf{k}_{\lambda_n}\|^2 \asymp e^{\psi(y_n) - 2y_n} (\psi''(y_n))^{1/2} = e^{h(\lambda_n)} / (\lambda_n \rho(\lambda_n)), \quad n \geq 0.$$

Proof. First of all, by Lemma 3.1, we have

$$\|\mathbf{k}_{\lambda_n}\|^2 \geq \frac{\lambda_n^{2n}}{\|z^n\|^2} \geq C e^{\psi(y_n) - 2y_n} (\psi''(y_n))^{1/2}.$$

Let $f \in \mathcal{F}_h$,

$$F(t) = \left(\int_0^{2\pi} |f(te^{i\theta})|^2 d\theta \right)^{1/2},$$

$$\omega(s) = \log F(\exp s).$$

Hardy's convexity theorem [6, Chapter 1] states that ω is convex.

Next, by (3.2), for large n we have $y_n - \alpha_n^{-1/2} > y_{n-1}$ and

$$\begin{aligned} \|f\|^2 &= \int_0^\infty F(t)^2 e^{-h(t)} t dt \geq \int_0^\infty e^{2\omega(s) - \psi(s) + 2s} ds \\ &\geq \int_{y_n - \alpha_n^{-1/2}}^{y_n + \alpha_n^{-1/2}} e^{2\omega(s) - \psi(s) + 2s} ds \\ &\geq C e^{-g(0)} \int_{-\alpha_n^{-1/2}}^{\alpha_n^{-1/2}} e^{2\omega(y_n + s) - 2n(y_n + s)} ds = I. \end{aligned}$$

If $\delta = \alpha_0^{-1/2}$, then by convexity of ω we have

$$\begin{aligned} I &\geq C e^{-g(0)} \alpha_n^{-1/2} \int_{-\delta}^\delta e^{2\omega(y_n + s) - 2n(y_n + s)} ds \\ &= C e^{-g(0)} \alpha_n^{-1/2} \int_{y_n - \delta}^{y_n + \delta} e^{2\omega(s) - 2ns} ds \\ &= C e^{-g(0)} \alpha_n^{-1/2} \int_{\exp(y_n - \delta)}^{\exp(y_n + \delta)} \frac{|f(z)|^2}{|z|^{2n+2}} dm(z). \end{aligned}$$

Applying the mean value theorem to the function $f(z)z^{-n-1}$, we conclude that

$$|f(e^{y_n})|^2 \leq C \|f\|^2 e^{g(0)} \alpha_n^{1/2} e^{2ny_n} = C \|f\|^2 e^{\psi(y_n) - 2y_n} \alpha_n^{1/2}.$$

□

Next, we consider a continuous piecewise linear function ℓ such that

$$\ell'(t) = 2n + 2, \quad y_n < t < y_{n+1}, \quad n \geq 0.$$

Then $\ell'(t) \leq \psi'(t)$ for large t , and hence, $\ell(t) \leq \psi(t) + O(1)$, $t \rightarrow \infty$. Next, $\ell'(t) + 2 \geq \psi'(t)$ for large t , and hence, $\ell(t) + 2t \geq \psi(t) + O(1)$, $t \rightarrow \infty$.

Lemma 3.3. *Given a small $\delta > 0$ we have*

- (a) $\int_0^\infty e^{\ell(t) - \psi(t)} dt < \infty$,
- (b) $\int_{y_n - \delta}^\infty e^{\ell(t) - \psi(t)} dt \leq C e^{\ell(y_n) - \psi(y_n)} (\psi''(y_n))^{-1/2}$,
- (c) $\int_0^{y_n - \delta} e^{\ell(t) - \psi(t) + 2t} dt \leq C e^{\ell(y_n) - \psi(y_n) + 2y_n} (\psi''(y_n))^{-1/2}$,
- (d) $\sum_{s=0}^n e^{\psi(y_s) - \ell(y_s)} (\psi''(y_s))^{1/2} \asymp e^{\psi(y_n) - \ell(y_n)} (\psi''(y_n))^{1/2}$,
- (e) $\sum_{s=n}^\infty e^{\psi(y_s) - \ell(y_s) - 2y_s} (\psi''(y_s))^{1/2} \asymp e^{\psi(y_n) - \ell(y_n) - 2y_n} (\psi''(y_n))^{1/2}$.

Proof. Set $u = \psi - \ell$. Then

$$\begin{aligned} u'(y_n + t) &\geq C\alpha_n t, & 0 \leq t \leq C\alpha_n^{-2/3}, \\ u'(y_n + t) &\geq C\alpha_n^{1/3}, & C\alpha_n^{-2/3} < t < y_{n+1} - y_n, \\ u'(y_n - t) &\geq 1, & 0 \leq t \leq C\alpha_n^{-2/3}. \end{aligned}$$

Therefore,

$$u(y_n) \geq u(y_{n-1}) + C\alpha_n^{-2/3}.$$

Since ψ'' does not increase, (a), (b), and (d) follow immediately.

Next, set $w(t) = \ell(t) + 2t - \psi(t)$. Then

$$\begin{aligned} w'(y_n + t) &\geq 1, & 0 \leq t \leq C\alpha_n^{-2/3}, \\ w'(y_n - t) &\geq C\alpha_n t, & 0 \leq t \leq C\alpha_n^{-2/3}, \\ w'(y_n - t) &\geq C\alpha_n^{1/3}, & C\alpha_n^{-2/3} < t < y_n - y_{n-1}. \end{aligned}$$

Therefore,

$$w(y_{n+1}) \geq w(y_n) + C\alpha_n^{-2/3}.$$

Since ψ'' does not increase, (c) and (e) follow immediately. □

Set

$$E(z) = \prod_{n \geq 0} \left(1 - \frac{z}{\lambda_n}\right).$$

Arguing like in the proof of Lemma 2.2, for small $\delta > 0$ we obtain

$$\begin{aligned} |E(e^t)|^2 &\leq C e^{\ell(t)}, \quad t \geq 0, \\ |E(e^t)|^2 &\asymp e^{\ell(t)}, \quad \text{dist}(t, \{y_n\}_{n \geq 0}) \geq \delta. \end{aligned}$$

Therefore, for every entire function $F \not\equiv 0$ we have

$$(3.4) \quad FE \notin \mathcal{F}_h.$$

Indeed,

$$\int_{\mathbb{C}} |F(z)|^2 |E(z)|^2 e^{-h(z)} dm(z) \geq C \int_0^\infty e^{\ell(t) - \psi(t) + 2t} dt = \infty.$$

In a similar way, by Lemma 3.3 (a), $E/(\cdot - \lambda_0) \in \mathcal{F}_h$. Furthermore,

$$(3.5) \quad |E'(\lambda_n)|^2 \asymp e^{\ell(y_n) - 2y_n}.$$

Set $d\mu(z) = |E(z)|^2 e^{-h(z)} dm(z)$, $v_n = e^{\psi(y_n) - \ell(y_n)} (\psi''(y_n))^{1/2}$, $\Omega_n = \{z : e^{y_n - \delta} < |z| < e^{y_{n+1} - \delta}\}$, $n \geq 0$. By Lemma 3.3 (b), we have

$$\begin{aligned} (3.6) \quad \int_{\Omega_n} \frac{d\mu(z)}{|z - \lambda_n|^2} &= \int_{e^{y_n - \delta} < |z| < e^{y_{n+1} - \delta}} \left| \frac{E(z)}{z - e^{y_n}} \right|^2 e^{-h(z)} dm(z) \\ &\asymp \int_{y_n - \delta}^{y_{n+1} - \delta} e^{\ell(t) - 2(t - y_n) - \psi(t) + 2(t - y_n)} dt \leq \int_{y_n - \delta}^\infty e^{\ell(t) - \psi(t)} dt \\ &\leq C \frac{e^{\ell(y_n) - \psi(y_n)}}{(\psi''(y_n))^{1/2}} = \frac{C}{v_n}, \quad n \geq 0. \end{aligned}$$

Furthermore, again by Lemma 3.3 (b), we have

$$\sum_{m=n+1}^\infty \int_{\Omega_m} \frac{d\mu(z)}{|z|^2} = \int_{y_{n+1} - \delta}^\infty e^{\ell(t) - \psi(t)} dt \leq \frac{C}{v_{n+1}}, \quad n \geq 0.$$

By Lemma 3.3 (d), we have

$$\sum_{s=0}^n v_s \asymp v_n, \quad n \geq 0.$$

Therefore,

$$(3.7) \quad \sum_{s=0}^n v_s \sum_{m=n+1}^\infty \int_{\Omega_m} \frac{d\mu(z)}{|z|^2} \leq C, \quad n \geq 0.$$

Next, by Lemma 3.3 (c), we have

$$\sum_{m=0}^n \int_{\Omega_m} d\mu(z) = \int_0^{y_{n+1} - \delta} e^{\ell(t) - \psi(t) + 2t} dt \leq C \frac{e^{2y_{n+1}}}{v_{n+1}}, \quad n \geq 0.$$

By Lemma 3.3 (e), we have

$$\sum_{s=n+1}^\infty \frac{v_s}{|\lambda_s|^2} \asymp \frac{v_{n+1}}{|\lambda_{n+1}|^2}, \quad n \geq 0.$$

Therefore,

$$(3.8) \quad \sum_{s=n+1}^\infty \frac{v_s}{|\lambda_s|^2} \sum_{m=0}^n \int_{\Omega_m} d\mu(z) \leq C, \quad n \geq 0.$$

Set $\Lambda = \{\lambda_n\}_{n \geq 0}$ and consider the mappings T_Λ and R_Λ defined in Subsection 1.1. The argument in the proof of Lemma 3.2 implies that the mapping T_Λ is bounded from \mathcal{F}_h to $\ell^2(1/\|\mathbf{k}_\lambda\|^2)$. By (3.4), T_Λ is injective. By Lemma 3.2 and by (3.5), $v_n \asymp \|\mathbf{k}_{\lambda_n}\|^2/|E'(\lambda_n)|^2$. Estimates (3.6)–(3.8) permit us to apply Theorem 1.1 in [2] to show that $d\mu$ is a Carleson measure for the space of the corresponding discrete Hilbert transforms. Thus, R_Λ is bounded from $\ell^2(1/\|\mathbf{k}_\lambda\|^2)$ to \mathcal{F}_h . By the argument in Subsection 1.1 we conclude that $\{\mathbf{k}_{\lambda_n}\}_{n \geq 0}$ is a Riesz basis in \mathcal{F}_h .

4. PROOF OF THEOREM 1.3

In this section we construct an increasing weight h and a corresponding set Λ consisting of (an increasing amount of) equidistributed points on rapidly growing concentric circles such that $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis in \mathcal{F}_h .

Given an increasing sequence $\{R_n\}_{n \geq 1}$ we set

$$\psi(t) = t + 2 \sum_{s \leq n} s(t - \log R_s) + n \min(t - \log R_n, \log R_{n+1} - t),$$

$$\log R_n \leq t \leq \log R_{n+1}, \quad n \geq 1.$$

Clearly, ψ is an increasing function.

An elementary geometric argument permits us to choose a sequence $\{R_n\}_{n \geq 1}$ satisfying the following two properties:

$$(I) \quad R_1 \geq 2, \quad R_{n+1} \geq R_n^2, \quad n \geq 1$$

and

$$(II) \quad \psi(t) = o(\varphi(t)), \quad t \rightarrow \infty.$$

In particular, $\log R_n \geq 2^n \log 2$. As above, we fix $h(t) = \psi(\log t)$. Next, we define

$$E(z) = \prod_{n \geq 1} \left(1 - \left(\frac{z}{R_n}\right)^n\right), \quad \Lambda = Z(E).$$

By an argument similar to that in the proof of Lemma 2.2, we have

$$(4.1) \quad |E(z)| \asymp e^{h(z)/2} \frac{n \operatorname{dist}(z, \Lambda)}{R_n^{3/2}}, \quad ||z| - R_n| \leq \frac{R_n}{n},$$

$$|E'(\lambda)| \asymp e^{h(R_n)/2} \frac{n}{R_n^{3/2}}, \quad \lambda \in \Lambda, |\lambda| = R_n,$$

$$|E(z)| \asymp e^{h(z)/2} |z|^{-1/2} \left|\frac{R_n}{z}\right|^{n/2}, \quad \frac{n+1}{n} R_n \leq |z| \leq \sqrt{R_n R_{n+1}},$$

$$|E(z)| \asymp e^{h(z)/2} |z|^{-1/2} \left|\frac{z}{R_{n+1}}\right|^{n/2}, \quad \sqrt{R_n R_{n+1}} \leq |z| \leq \frac{n R_{n+1}}{n+1}.$$

Let $F \neq 0$ be an entire function. Then

$$(4.2) \quad \int_{\mathbb{C}} |F(z)|^2 |E(z)|^2 e^{-h(z)} dm(z)$$

$$\geq C \sum_{n \geq 1} \int_{\frac{n-1}{n} \leq \frac{|z|}{R_n} \leq \frac{n+1}{n}} |F(z)|^2 \frac{n^2 \operatorname{dist}^2(z, \Lambda)}{R_n^3} dm(z) \geq \sum_{n \geq 1} \frac{R_n}{n} = \infty.$$

Thus, $E \notin \mathcal{F}_h$ and for any entire function $F \neq 0$ we have $FE \notin \mathcal{F}_h$.

Given $\lambda \in \Lambda$, we set $E_\lambda = \frac{E}{\cdot - \lambda}$. Next, we are going to deal with the scalar products $\langle E_\lambda, E_\mu \rangle_{\mathcal{F}_h}$. For $\lambda \in \Lambda$, $|\lambda| = R_n$, we have

$$\begin{aligned}
(4.3) \quad \|E_\lambda\|^2 &= \int_{\mathbb{C}} \frac{|E(z)|^2}{|z - \lambda|^2} e^{-h(z)} dm(z) \\
&\leq C \left(\int_{\frac{n-1}{n} \leq \frac{|z|}{R_n} \leq \frac{n+1}{n}} \frac{n^2 \text{dist}^2(z, \Lambda)}{R_n^3 |z - \lambda|^2} dm(z) \right. \\
&\quad + \sum_{s \geq 1, s \neq n} \int_{\frac{s-1}{s} \leq \frac{|z|}{R_s} \leq \frac{s+1}{s}} \frac{s^2 \text{dist}^2(z, \Lambda)}{R_s^3 |z - \lambda|^2} dm(z) \\
&\quad + \sum_{s \geq 1} \int_{\frac{s+1}{s} R_s \leq |z| \leq \sqrt{R_s R_{s+1}}} \left| \frac{R_s}{z} \right|^s \frac{dm(z)}{|z| \cdot |z - \lambda|^2} \\
&\quad + \sum_{s \geq 1} \int_{\sqrt{R_s R_{s+1}} \leq |z| \leq \frac{s}{s+1} R_{s+1}} \left| \frac{z}{R_{s+1}} \right|^s \frac{dm(z)}{|z| \cdot |z - \lambda|^2} \Bigg) \\
&\leq \frac{C}{R_n} + C \sum_{s \geq 1, s \neq n} \frac{R_s}{s \max(R_s, R_n)^2} \asymp \frac{1}{R_n}.
\end{aligned}$$

Furthermore, if $\lambda, \mu \in \Lambda$, $R_j = |\lambda| < |\mu| = R_k$, then in a similar way

$$(4.4) \quad \int_{\mathbb{C}} \frac{|E(z)|^2}{|z - \lambda| \cdot |z - \mu|} e^{-h(z)} dm(z) \leq \frac{C \log(k+1)}{R_k}.$$

Next we fix $n \geq 1$ and set $R = R_n$. Then

$$E(z) = \left(1 + O\left(\frac{1}{R}\right)\right) \left(1 - \left(\frac{z}{R}\right)^n\right) \prod_{1 \leq s < n} \left(\frac{z}{R_s}\right)^s, \quad ||z| - R| \leq \frac{R}{2}.$$

Let $\alpha \neq \beta$ be such that $|\alpha| = |\beta| = 1$, and let $\lambda = \alpha R$, $\mu = \beta R$ belong to Λ . We have

$$\begin{aligned}
&\int_{\mathbb{C}} \frac{|E(z)|^2}{(z - \lambda)(\overline{z - \mu})} e^{-h(z)} dm(z) \\
&= O\left(\frac{1}{nR}\right) + \int_{R/2 \leq |z| \leq R} \frac{\left(1 - \left(\frac{z}{R}\right)^n\right) \left(1 - \left(\frac{\bar{z}}{R}\right)^n\right)}{(z - \lambda)(\overline{z - \mu})} \left|\frac{z}{R}\right|^{n-1} \frac{dm(z)}{|z|} \\
&\quad + \int_{R \leq |z| \leq 3R/2} \frac{\left(1 - \left(\frac{z}{R}\right)^n\right) \left(1 - \left(\frac{\bar{z}}{R}\right)^n\right)}{(z - \lambda)(\overline{z - \mu})} \left|\frac{R}{z}\right|^{3n} \frac{dm(z)}{|z|} \\
&= O\left(\frac{1}{nR}\right) + \frac{1}{R} \int_{1/2}^1 \left(\sum_{s=0}^{n-1} \alpha^s w^{n-1-s}\right) \overline{\left(\sum_{s=0}^{n-1} \beta^s w^{n-1-s}\right)} |w|^{n-2} dm(w) \\
&\quad + \frac{1}{R} \int_1^{3/2} \left(\sum_{s=0}^{n-1} \alpha^s w^{n-1-s}\right) \overline{\left(\sum_{s=0}^{n-1} \beta^s w^{n-1-s}\right)} |w|^{-3n-1} dm(w) \\
&= O\left(\frac{1}{nR}\right) + \frac{1}{R} \sum_{s=0}^{n-1} 2\pi (\alpha \bar{\beta})^s \left(\frac{1}{2s + n + 1} + \frac{1}{3n - 2s - 2}\right).
\end{aligned}$$

Set

$$a_s = \frac{1}{2s + n + 1} + \frac{1}{3n - 2s - 2}, \quad 0 \leq s < n,$$

$$b_{j,k} = \sum_{s=0}^{n-1} e^{2\pi i(j-k)s/n} a_s, \quad 0 \leq j, k < n,$$

$$A = (b_{j,k})_{0 \leq j, k < n}.$$

Then A is a circulant matrix. It is well-known and easily verified that A has an orthonormal basis of eigenvectors $n^{-1/2}(e^{2\pi i k q/n})_{0 \leq k < n}$, $0 \leq q < n$, and the corresponding eigenvalues are $\sum_{k=0}^{n-1} b_{0,k} e^{2\pi i k q/n}$, $0 \leq q < n$. Therefore, the norm of the corresponding operator acting on ℓ_n^2 is equal to

$$\max_{0 \leq q < n} \left| \sum_{k=0}^{n-1} e^{2\pi i k q/n} \sum_{s=0}^{n-1} e^{-2\pi i k s/n} a_s \right| = n \max_{0 \leq q < n} |a_q| = O(1), \quad n \rightarrow \infty.$$

Thus, if

$$B_n = \left(\langle E_\lambda, E_\mu \rangle_{\mathcal{F}_h} \right)_{\lambda, \mu \in \Lambda, |\lambda|=|\mu|=R_n},$$

then

$$(4.5) \quad \|B_n\|_{\ell_n^2 \rightarrow \ell_n^2} = O\left(\frac{1}{R_n}\right), \quad n \rightarrow \infty.$$

Now, for $|\lambda| = R_n$ we estimate $\|\mathbf{k}_\lambda\|$. Applying the mean value theorem to the function

$$g(z) = f(z) \prod_{1 \leq s < R_n} \left(\frac{R_s}{z}\right)^s$$

in the disc $D = \{z : |z - \lambda| < R_n/n\}$, we obtain that

$$(4.6) \quad |f(\lambda)|^2 e^{-h(\lambda)} \leq \frac{Cn^2}{R_n^2} \int_D |f(w)|^2 e^{-h(w)} dm(w),$$

and hence,

$$\|\mathbf{k}_\lambda\| \leq C e^{h(R_n)/2} \frac{n}{R_n}, \quad |\lambda| = R_n.$$

By (4.1) and (4.3), for $\lambda \in \Lambda$ such that $|\lambda| = R_n$ we have

$$\|\mathbf{k}_\lambda\| \geq \frac{|E_\lambda(\lambda)|}{\|E_\lambda\|} = \frac{|E'(\lambda)|}{\|E_\lambda\|} \geq C e^{h(R_n)/2} \frac{n}{R_n}.$$

Replacing $E(z)$ by $E(e^{i\theta}z)$ we get the same estimate for all λ such that $|\lambda| = R_n$. Thus,

$$(4.7) \quad \frac{1}{C} e^{h(R_n)/2} \frac{n}{R_n} \leq \|\mathbf{k}_\lambda\| \leq C e^{h(R_n)/2} \frac{n}{R_n}, \quad |\lambda| = R_n,$$

for some C independent of λ and n .

Now, we consider the mappings T_Λ and R_Λ defined in Subsection 1.1. Estimate (4.6) implies that the mapping T_Λ is bounded from \mathcal{F}_h to $\ell^2(1/\|\mathbf{k}_\lambda\|^2)$. By (4.2), T_Λ is injective. Set $e_\lambda = \{\|\mathbf{k}_\lambda\| \delta_{\lambda\mu}\}_{\mu \in \Lambda}$. Then $\|e_\lambda\|_{\ell^2(1/\|\mathbf{k}_\lambda\|^2)} = 1$ and $R_\Lambda e_\lambda = E_\lambda \|\mathbf{k}_\lambda\| / E'(\lambda)$. Estimates (4.1), (4.4), (4.5), and (4.7) show that the matrix $(\langle R_\Lambda e_\lambda, R_\Lambda e_\mu \rangle)_{\lambda, \mu \in \Lambda}$ consists of (i) a sequence of squares along the diagonal with uniformly bounded norms and (ii) rapidly decaying

off-diagonal terms. Thus, R_Λ is bounded from $\ell^2(1/\|\mathbf{k}_\lambda\|^2)$ to \mathcal{F}_h . By the argument in Subsection 1.1 we conclude that $\{\mathbf{k}_\lambda\}_{\lambda \in \Lambda}$ is a Riesz basis in \mathcal{F}_h .

Finally, for $n \geq 3$ we set

$$f_n(z) = \frac{E(z)}{z - R_n} \prod_{1 \leq s < n/2} \frac{z - e^{-2\pi i s/n} R_n}{z - e^{2\pi i s/n} R_n}.$$

Then $|E_{R_n}| = |f_n|$ on the real line, and

$$\begin{aligned} \|f_n\|^2 &= \int_{\mathbb{C}} |f_n(z)|^2 e^{-h(z)} dm(z) \\ &\geq \int_{\{|z - R_n| \leq \frac{R_n}{n}, |\arg z - \pi/2| \leq \pi/4\}} |f_n(z)|^2 e^{-h(z)} dm(z) \\ &\geq \frac{C}{nR_n} 2^{n/20} \geq n \|E_{R_n}\|^2, \quad n \rightarrow \infty. \end{aligned}$$

This shows that \mathcal{F}_h is not (up to the norm equivalence) isomorphic to a de Branges space.

5. PROOF OF PROPOSITION 1.4

Let $\mathcal{F}_h = \mathcal{F}_{h_0}$ (up to the norm equivalence), and let the polynomials belong to \mathcal{F}_{h_0} . We have

$$\|z^n\|_{\mathcal{F}_h}^2 = \int_0^\infty 2\pi r^{2n+1} e^{-h(r)} dr = \int_0^\infty x^n u_h(x) dx,$$

where $u_h(x) = \pi e^{-h(\sqrt{x})}$.

If h is continuous and non-decreasing, we can find a non-increasing $\tilde{u} \in C^1(\mathbb{R})$ such that $u_h \leq \tilde{u} \leq 2u_h$, $\tilde{u}' \leq 0$. Since

$$\int_0^\infty x^n \tilde{u}(x) dx = -\frac{1}{n+1} \int_0^\infty x^{n+1} \tilde{u}'(x) dx,$$

by the Cauchy–Schwarz inequality we obtain that for some bounded sequence $\{c_n\}_{n \geq 1}$,

$$(5.1) \quad \text{the sequence } \{\log[n\|z^n\|_{\mathcal{F}_{h_0}}^2] + c_n\}_{n \geq 1} \text{ is convex.}$$

On the other hand, let

$$p_n = \log \int_0^\infty x^n d\mu(x),$$

where

$$d\mu(x) = \sum_{k \geq 1} e^{-s_k t_k} \delta_{\exp s_k},$$

$t_k \in \mathbb{N}$, $s_k \geq 1$, $k \geq 0$. If $t_k > t_{k-1} + 1$, $s_k > 2t_k s_{k-1}$, $k > 1$, then

$$p_n = \log \sum_{k \geq 1} e^{(n-t_k)s_k} = \max_{k \geq 1} [(n-t_k)s_k] + O(1), \quad n \geq 1,$$

and

$$p_n = (n-t_k)s_k + O(1)$$

for $t_k + 1 \leq n \leq t_{k+1}$, $k \geq 1$.

Finally, let $t_k \geq t_{k-1}^2$, $k > 1$, let u_{h_0} be a continuous positive function such that

$$\left| p_n - \log \int_0^\infty x^n u_{h_0}(x) dx \right| \leq 1, \quad n \geq 1,$$

and let $u_{h_0}(x) = \pi e^{-h_0(\sqrt{x})}$. Then h_0 is continuous and there exists a bounded sequence $\{d_n\}_{n \geq 1}$ such that the sequence

$$\{\log[\|z^n\|_{\mathcal{F}_{h_0}}^2 + d_n]\}_{n \geq 1}$$

is linear for $t_k + 1 \leq n \leq t_{k+1}$, $k \geq 1$, which contradicts to (5.1).

REFERENCES

- [1] A. Baranov, Y. Belov, A. Borichev, *Spectral synthesis in de Branges spaces*, Geometric and Functional Analysis **25** (2015), 417–452.
- [2] Yu. Belov, T. Mengestie, K. Seip, *Discrete Hilbert transforms on sparse sequences*, Proc. London Math. Soc. **103** (2011), 73–105.
- [3] A. Borichev, R. Dhuez, K. Kellay, *Sampling and interpolation in large Bergman and Fock spaces*, Journal of Functional Analysis **242** (2007), 563–606.
- [4] A. Borichev, Yu. Lyubarskii, *Riesz bases of reproducing kernels in Fock type spaces*, J. Inst. Math. Jussieu **9** (2010), 449–461.
- [5] L. de Branges, *Hilbert Spaces of Entire Functions*, Prentice–Hall, Englewood Cliffs, 1968.
- [6] P. Duren, *Theory of H^p spaces*, Pure and Applied Mathematics, Vol. 38, Academic Press, New York–London, 1970.
- [7] W. Hayman, *The local growth of power series: a survey of the Wiman–Valiron method*, Canad. Math. Bull. **17** (1974), 317–358.
- [8] Yu. Lyubarskii, E. Malinnikova, *On approximation of subharmonic functions*, J. Anal. Math. **83** (2001), 121–149.
- [9] N. Nikolski, *Operators, Functions, and Systems: an Easy Reading. Vol. 2*, Math. Surveys Monogr., Vol. 93, AMS, Providence, RI, 2002.
- [10] K. Seip, *Density theorems for sampling and interpolation in the Bargmann–Fock space*, I, J. Reine Angew. Math. **429** (1992), 91–106.

ANTON BARANOV,

DEPARTMENT OF MATHEMATICS AND MECHANICS, ST. PETERSBURG STATE UNIVERSITY,
ST. PETERSBURG, RUSSIA,
NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, ST. PETERSBURG,
RUSSIA,
anton.d.baranov@gmail.com

YURI BELOV,

CHEBYSHEV LABORATORY, ST. PETERSBURG STATE UNIVERSITY, ST. PETERSBURG,
RUSSIA,
j_b_juri_belov@mail.ru

ALEXANDER BORICHEV,

I2M, CNRS, CENTRALE MARSEILLE, AIX-MARSEILLE UNIVERSITÉ, 13453 MARSEILLE,
FRANCE,
DEPARTMENT OF MATHEMATICS AND MECHANICS, SAINT PETERSBURG STATE UNIVER-
SITY, ST. PETERSBURG, RUSSIA,
alexander.borichev@math.cnrs.fr